

#42, 100FT ROAD, KAMMAGONDANAHALLI, JALAHALLI WEST, **BENGALURU - 560 015** Phone: 9008030463 9008030896 / 9513330437


HOSKOTE - MALUR ROAD, ISRI CROSS, KATTIGENAHALLI, JADIGENAHALLI HOBLI, **BENGALURU - 562114** Phone: 9741332998 7760575487/9535527713

PLOT NO.87, VAHINI NIVAS MATRUSRI NAGAR COLONY, HAFEEZ PET, MIYAPUR, **HYDERABAD - 500049** Phone: 8977548407 8977548408 / 8977548409

KCET EXAMINATION - 2024

VERSION: A2 SUBJECT: CHEMISTRY

DATE: 19-04-2024





- 1. Which of the following set of polymers are used as fibre?
 - (i) Teflon
 - (ii) Starch
 - (iii) Terylene
 - (iv) Orlon
 - (A) (i) and (ii)
 - (B) (ii) and (iii)
 - (C) (iii) and (iv)
 - (D) (i) and (iv)

Ans. C


Sol. Terylene and orlon are fibres

- 2. The biodegradable polymer obtained by polymerisation, of Glycine and Aminocaproic acid is
 - (A) Nylon 6
 - (B) PHBV
 - (C) Nylon 2 Nylon 6
 - (D) Nylon 6, 10

Ans. C

Sol. Nylon 2 – Nylon 6 is a biodegradable polymer

3. The compound



- (A) Sucralose
- (B) Aspartame
- (C) Saccharin
- (D) Alitame

Ans. NO OPTION

Sol. Saccharin structure

- 4. Which one of the following is a cationic detergent?
 - (A) Cetyltrimethylammonium bromide
 - (B) Sodium dodecylbenzene sulphonate
 - (C) Dodecylbenzene sulphonic acid
 - (D) Dodecylbenzene

Ans. A

Sol. Cetyltrimethylammonium bromide is a cationic detergent

- 5. The type of linkage present between nucleotides is
 - (A) Phosphoester linkage
 - (B) Phosphodiester linkage
 - (C) Amide linkage
 - (D) Glycosidic linkage

Ans. B

- **Sol.** Nucleotides are joined by Phosphodiester linkage
- 6. $\alpha D (+) \text{ glucose and } \beta D (+) \text{ glucose}$ are
 - (A) Enantiomers
 - (B) Conformers
 - (C) Epimers
 - (D) Anomers

Ans. D

- **Sol.** $\alpha D (+) -$ glucose and $\beta D (+) -$ glucose are anomeric at first carbon so called anomers
- 7. Propanone and Propanal are
 - (A) Position isomers
 - (B) Functional isomers
 - (C) Chain isomers
 - (D) Geometrical isomers

Ans. B

- Sol. Propanone and Propanal are functional isomers
- 8. Sodium ethanoate on heating with soda lime gives 'X'. Electrolysis of aqueous solution of sodium ethanoate gives 'Y'. 'X' and 'Y' respectively are
 - (A) Methane and Ethane
 - (B) Methane and Methane
 - (C) Ethane and Methane
 - (D) Ethane and Ethane


Ans. A

Sol. $CH_3COONa + NaOH \xrightarrow{CaO} CH_4 + Na_2CO_3$ X $2CH_3COONa + 2H_2O \xrightarrow{\Delta} C_2H_6 + 2NaOH + 2CO_2 + H_2$ Y

 $X = CH_4$ $Y = C_2H_6$



9. But-1-yne on reaction with $dil.H_2SO_4$ in presence of Hg^{2+} ions at 333K gives



Ans. A Sol.

$$CH_{3} - CH_{2} - C \equiv CH + H_{2}O \xrightarrow{Hg^{+2}} CH_{3} - CH_{2} - C = CH_{2}$$

$$OH$$

$$Tautamerises$$

$$CH_{3} - CH_{2} - C - CH_{3}$$

$$CH_{3} - CH_{2} - C - CH_{3}$$

$$(or)$$

- 10. Biologically active adrenaline and ephedrine used to increase blood pressure contain
 - (A) Primary amino group
 - (B) Secondary amino group
 - (C) Tertiary amino group
 - (D) Quaternary ammonium salt

Ans. B

Sol. Adrenaline and ephedrine contains secondary amino group

11. In the reaction

Aniline
$$\xrightarrow{\text{NaNO}_2} P \xrightarrow{\text{Phenol}} Q$$
.

- (A) $C_6H_5N_2C1$
- (B) ortho-hydroxyazobenzene
- (C) para-hydroxyazobenzene
- (D) meta-hydroxyazobenzene


Ans. C Sol.

- 12. The female sex hormone which is responsible for the development of secondary female characteristics and participates in the control of menstrual cycle is
 - (A) Testosterone
- (B) Estradiol
- (C) Insulin
- (D) Thyroxine

Ans. B

Sol. Conceptual

13. In the following scheme of reaction.




X, Y and Z respectively are:

- (A) AgF, alcoholic KOH and benzene
- (B) HF, aqueous KOH and Na in dry ether
- (C) Hg₂F₂, alcoholic KOH and Na in dry ether
- (D) CoF₂, aqueous KOH and benzene

Ans. C

Sol.



- 14. 8.8 g of monohydric alcohol added to ethyl magnesium iodide in ether liberates 2240 cm³ of ethane at STP. This monohydric alcohol when oxidised using pyridinium-chloromate, forms a carbonyl compound that answers silver mirror test (Tollen's test). The monohydric alcohol is
 - (A) butan-2-ol
 - (B) 2, 2-dimethyl propan-1-ol
 - (C) pentan-2-ol
 - (D) 2, 2-dimethyl ethan-1-ol

Ans. B

Sol. 8.8 g of monohydric alcohol \leftarrow 2240 cm³

88 g
$$\leftarrow 22400 \text{ cm}^3$$

The carbonyl compound that can oxidise tollens reagent i.e., **Aldehyde** and which is oxidising product of Primary alcohol.



- 15. When a tertiary alcohol 'A' $(C_4H_{10}O)$ reacts with 20% H_3PO_4 at 358 K, it gives a compound 'B' (C_4H_8) as a major product. The IUPAC name of the compound 'B' is
 - (A) But-1-ene
 - (B) But-2-ene
 - (C) Cyclobutane
 - (D) 2-Methylpropene

Ans. D Sol.

- 16. PCC is
 - (A) $K_2Cr_2O_7 + Pyridine$
 - (B) $CrO_3 + CHCl_3$
 - (C) $CrO_3 + H_2SO_4$
 - (D) A complex of chromium trioxide with pyridine + HCl

Ans. D

- **Sol.** PCC is pyridinium chlorochromate C_5H_5 NHCrO₃Cl⁻
- 17. On treating 100 mL of 0.1 M aqueous solution of the complex CrCl₃.6H₂O with excess of AgNO₃, 2.86 g of AgCl was obtained. The complex is
 - (A) $[Cr(H_2O)_3Cl_3].3H_2O$
 - (B) [Cr(H₂O)₄Cl₂]Cl.2H₂O
 - (C) $[Cr(H_2O)_5Cl]Cl_2.H_2O$
 - (D) $[Cr(H_2O)_6Cl_3]$

Ans. C

Sol. 0.01 moles of an aqueous solution of $CrCl_3.6H_2O$

given 2.86 g AgCl means,

1 mole of aqueous solution = 2 moles of AgCl

 $143.5 \times 2 = 286g \text{ of AgCl}$ So, $[Cr(H_2O)_5Cl]Cl_2.H_2O$

- 18. The complex compounds $[Co(NH_3)_5SO_4]Br$ and $[Co(NH_3)_5Br]SO_4$ are
 - (A) Coordination isomers
 - (B) Geometrical isomers
 - (C) Optical isomers
 - (D) Ionisation isomers

Ans. D

- **Sol.** $[Co(NH_3)_5SO_4]Br$ and $[Co(NH_3)_5Br]SO_4$ are Ionisation isomers.
- 19. When of the following statements are true about $[CoF_6]^{3-}$ ion?
 - I) The complex has octahedral geometry.
 - II) Coordination number of Co is 3 and oxidation state is +6.
 - (III) The complex is sp³d² hybridised
 - (IV) It is a high spin complex
 - (A) I, II and IV
 - (B) I, III and IV
 - (C) II and IV
 - (D) II, III and IV

Ans. B

- Sol. I, III and IV statements are true.
- 20. A haloalkane undergoes S_{N^2} or S_{N^1} reaction depending on
 - (A) Solvent used in the reaction
 - (B) Low temperature
 - (C) The type of halogen atom
 - (D) Stability of the haloalkane

Ans. A

- **Sol.** S_{N^2} or S_{N^1} reaction depends on solvent used in the reaction.
- 21. 2-Methyl propane can be prepared by Wurtz reaction. The haloalkanes taken along with metallic sodium and dry ether are:
 - (A) chloromethane and 2-chloropropane
 - (B) chloroethane and chloromethane
 - (C) chloroethane and 1-chloropropane
 - (D) chloromethane and 1-chloropropane

Ans. A

Sol.

$$\begin{array}{c} \operatorname{CH_3-Cl+CH_3-CH-CH_3} \xrightarrow{\quad \text{CNa} \quad \text{dry ether} \quad} \operatorname{CH_3-CH-CH_3+2NaCl} \\ \operatorname{Chloromethane} & \operatorname{Cl} & \operatorname{CH_3} \\ \text{2- chloro propane} & \text{2-methyl propane} \end{array}$$



In the analysis of III group basic radicals of salts, the purpose of adding NH₄Cl_(s) to

NH₄OH is:

- (A) to increase the concentration of OH⁻ ions.
- (B) to precipitate the radicals of group IV and V .
- (C) to suppress the dissociation of NH₄OH.
- (D) to introduce Cl⁻ions.

Ans. C

- **Sol.** To suppress the dissociation of NH₄OH due to the common ion effect
- Solubility product of CaC2O4 at a given 23. temperature in pure water is $4 \times 10^{-9} (\text{moL}^{-1})^2$ Solubility of CaC₂O₄ at the same temperature
 - (A) $6.3 \times 10^{-5} \text{ mol L}^{-1}$ (B) $2 \times 10^{-5} \text{ mol L}^{-1}$
 - (C) $2 \times 10^{-4} \text{ mol L}^{-1}$
- (D) $6.3 \times 10^{-4} \text{ mol L}^{-1}$

Ans. A

 $CaC_2O_4 \to Ca^{2+} + C_2O_4^{2-}$ Sol.

$$K_{sp} = (S)(S)$$

$$4 \times 10^{-9} = S^2$$

$$S = \sqrt{40 \times 10^{-10}} = 6.3 \times 10^{-5} \text{ mol / L}$$

- 24. In the reaction between moist SO₂ acidified permanganate solution:
 - (A) SO_2 is oxidised to SO_4^{2-}

MnO₄ is reduced to Mn²⁺

(B) SO₂ is reduced to S

MnO₄ is oxidised to MnO₄

(C) SO_2 is oxidised to SO_3^{2-}

MnO₄ is reduced to MnO₂

(D) SO₂ is reduced to H₂S

MnO₄ is oxidised to MnO₄

Ans. A

Sol. $2MnO_4^- + 3SO_2 + 4H^+ \rightarrow 2Mn^{2+} + 3SO_4^{2-} + 2H_2O$

- 25. Which one of the following properties is generally not applicable to ionic hydrides?
 - (A) Non-volatile
 - (B) Non-conducting in solid state
 - (C) Crystalline
 - (D) Volatile

Ans. D

- Sol. Ionic hydrides exhibit high Mpt i.e, Non -
- 26. Which one of the following nitrate will decompose to give NO2 on heating?
 - (A) NaNO₃
 - (B) KNO₃
 - (C) RbNO₃
 - (D) LiNO₃

Ans. D

Sol.
$$4\text{LinO}_3 \xrightarrow{\Delta} 2\text{Li}_2\text{O} + 4\text{NO}_2 + \text{O}_2$$

- Which of the following halides cannot be 27. hydrolysed?
 - (A) CCl₄
 - (B) SiCl₄
 - (C) GeCl₄
 - (D) SnCl₄

Ans. A

- **Sol.** Due to the absence of d orbitals.
- 0.48g of an organic compound on complete 28. combustion produced 0.22 g of CO₂. The percentage of C in the given organic compound is:
 - (A) 25
 - (B) 50
 - (C) 12.5
 - (D) 87.5

Ans. C

Sol. % C =
$$\frac{12}{44} \times \frac{\text{wt. of CO}_2}{\text{wt. of O.C}} \times 100$$

%C =
$$\frac{12}{44} \times \frac{0.22}{0.48} \times 100 = \frac{600}{48} = 12.5$$
%



29. In the given sequence of reactions, identify 'P', 'Q' and 'S' respectively.

$$CH_2 = CH_2 \xrightarrow{P} CH_2 - CH_2 \xrightarrow{Q} Br Br$$

$$CH_2 = CH - Br \xrightarrow{R} CH \equiv CH \xrightarrow{S} C_6H_6$$

- (A) Br₂, Alc, KOH, NaOH, Al₂O₃
- (B) HBr, Alc, KOH, CaC₂, KMnO₄
- (C) HBr, Alc, KOH, NaNH2, Red hot iron tube
- (D) Br₂, Alc, KOH, NaNH₂, Red hot iron tube

Ans. D Sol

Sol
$$CH_{2} = CH_{2} \xrightarrow{Br_{2}} CH_{2} \xrightarrow{CH_{2}} CH_{2} \xrightarrow{Alc.KOH} CH_{2} = CH - Br \xrightarrow{NaNH_{2}} CH = CH$$

$$CH = CH$$

$$CH = CH$$

- 30. The first chlorinated organic insecticide propared is:
 - (A)Gammaxene
 - (B)Chloroform
 - (C) COCl₂
 - (D)DDT

Ans. D

Sol. Dichloro Diphenyl Trichloro ethane (D.D.T)

- 31. Which of the following crystals has the unit cell such that $a = b \neq c$ and $\alpha = \beta = 90^{\circ}$, $\gamma = 120^{\circ}$?
 - (A) Zinc blende
 - (B) Graphite
 - (C) Cinnabar
 - (D) Potassium dichromate

Ans. B

Sol. Conceptual

- 32. MnO exhibits:
 - (A) Ferrimagnetism
 - (B) Antiferromagnetism
 - (C) Ferromagnetism
 - (D) Paramagnetism

Ans. B

Sol. Conceptual

33. The number of atoms in 4.5g of a face-centred cubic crystal with edge length 300pm is: (Given density = 10 g cm^{-3} and

$$N_A = 6.022 \times 10^{23})$$

- (A) 6.6×10^{20}
- (B) 6.6×10^{23}
- (C) 6.6×10^{19}
- (D) 6.6×10^{22}

Ans. D

Sol.
$$d = \frac{Z \times M}{N_{\Delta} \times a^3}$$

$$\Rightarrow M = \frac{10 \times 6.022 \times 10^{23} \times (300 \times 10^{-10})^{3}}{4}$$

$$M=40.5\,gm$$

Therefore 40.5 gm \rightarrow 6.022×10²³ atoms 4.5gm \rightarrow x

$$x = 6.6 \times 10^{22}$$
 atoms

- 34. Vapour pressure of a solution containing 18 g of glucose and 178.2 g of water at 100°C is:

 (Vapour pressure of pure water at 100°C = 760 torr)
 - (A) 76.0 torr
 - (B) 752.0 torr
 - (C) 7.6 torr
 - (D) 3207.6 torr

Ans. B

Sol.
$$n_{H_2O} = \frac{178.2}{18} = 9.9$$
; $n_{C_6H_{12}O_6} = \frac{18}{180} = 0.1$

$$\chi_{C_6H_{12}O_6} = \frac{0.1}{10} = 0.01$$

$$\frac{P_0 - P_s}{P_0} = \chi_{C_6 H_{12} O_6}$$

$$\frac{760 - P_s}{760} = 0.01$$

$$760 - P_s = 7.6$$

$$P_s = 752.4 torr$$

- 35. A mixture of phenol and aniline shows negative deviation from Raoult's law. This is due to the formation of:
 - (A) Polar covalent bond
 - (B) Non-polar covalent bond
 - (C) Intermolecular Hydrogen bond
 - (D) Intramolecular Hydrogen bond

Ans. C

Sol. Due to intermolecular hydrogen bond



- 36. Which one of the following pairs will show positive deviation from Raoult's Law?
 - (A) Water HCl
 - (B) Benzene-Methanol
 - (C) Water -HNO₃
 - (D) Acetone Chloroform

Ans. B

- **Sol.** Conceptual
- 37. How many Coulombs are required to oxidise $0.1 \text{ mole of } H_2O$ to oxygen?
 - (A) 1.93×10^5 C
 - (B) 1.93×10^4 C
 - (C) 3.86×10^4 C
 - (D) 9.65×10^{3} C

Ans. B

Sol.
$$H_2O \rightarrow H_2 + \frac{1}{2}O_2$$

1 mole $H_2O \rightarrow 2$ Faradays = $2 \times 96500C$

 $0.1 \text{ mole } H_2O \rightarrow xC$

$$x = \frac{2 \times 96500 \times 0.1}{1} = 19300C = 1.93 \times 10^{4}C$$

- 38. A current of 3A is passed through a molten calcium salt for 1hr 47 min 13sec. The mass of calcium deposited is:
 - (Molar mass of $Ca = 40 \text{ g mol}^{-1}$)
 - (A) 6.0 g
 - (B) 2.0 g
 - (C) 8.0 g
 - (D) 4.0 g

Ans. D

Sol.
$$w = \frac{Eit}{96500}$$

= $\frac{20 \times 3 \times 6432}{96500}$
= $3.99 \approx 4$

- 39. The value of 'A' in the equation $\frac{1}{2}$
 - $\lambda_{\rm m} = \lambda_{\rm m}^0 A\sqrt{C}$ is same for the pair
 - (A) NaCl and CaCl₂
 - (B) CaCl2 and MgSO4
 - (C) NaCl and KBr
 - (D) MgCl2 and NaCl

Ans. C

Sol. Conceptual

- 40. For the reaction, $A \rightleftharpoons B$, $E_a = 50 \text{ kJ mol}^{-1}$ and $\Delta H = -20 \text{ kJ mol}^{-1}$. When a catalyst is added, E_a decreases by 10 kJ mol $^{-1}$. What is the E_a for the backward reaction in the presence of catalyst?
 - (A) 60 kJ mol⁻¹
 - (B) 40 kJ mol⁻¹
 - (C) 70 kJ mol⁻¹
 - (D) 20 kJ mol⁻¹

Ans. A

Sol.
$$\Delta H = (E_a)_f - (E_a)_b$$

 $-20 = 40 - (E_a)_b$

$$\left(E_a\right)_b=60$$

41. For the reaction $PCl_5 \rightarrow PCl_3 + Cl_2$, rate and rate constant are $1.02 \times 10^{-4} \text{ mol L}^{-1}\text{S}^{-1}$ and

 3.4×10^{-5} s⁻¹ respectively at a given instant. The molar concentration of PCl₅ at that instant is:

- (A) 8.0 mol L^{-1}
- (B) $3.0 \text{ mol } L^{-1}$
- (C) $0.2 \text{ mol } L^{-1}$
- (D) $2.0 \text{ mol } L^{-1}$

Ans. B

Sol. Rate = $k[PCl_5]$

$$\frac{1.02 \times 10^{-4}}{3.4 \times 10^{-5}} = [PCl_5]$$

:: [PCl₅] = 3.0 mol / lit

42. Which one of the following does not represent Arrhenius equation?

(A)
$$\log k = \log A - \frac{Ea}{2.303RT}$$

- (B) $k = Ae^{-Ea/RT}$
- (C) In $k = -\frac{Ea}{RT} + In A$
- (D) $k = Ae^{Ea/RT}$

Ans. D

Sol. Conceptual



- 43. Identify the *incorrect* statement:
 - (A) Values of colligative properties of colloidal solution are of small order compared to values of true solution
 - (B) Tyndall effect is observed only when diameter of the dispersed particles is not much smaller than wavelength of incident light
 - (C) Colour of colloidal solution depends on the wavelength of light scattered by the dispersed particles
 - (D) Brownian movement is due to balanced bombardment of molecules of dispersion medium on colloidal particles

Ans. D

- **Sol.** Brownian movement is due to **unbalanced** bombardment of molecules of dispersion medium on colloidal particles
- 44. For the coagulations of positively charged hydrated ferric oxide sol, the flocculating power of the ions is in the order:

(A)
$$PO_4^{3-} > SO_4^{2-} > Cl^- > [Fe(CN)_6]^4$$

(B)
$$C1^- > SO_4^{2-} > PO_4^{3-} > \left[Fe(CN)_6 \right]^{4-}$$

(C)
$$SO_4^{2-} = Cl^- = PO_4^{3-} = \left[Fe(CN)_6 \right]^{4-}$$

(D)
$$\left[\text{Fe} \left(\text{CN} \right)_{6} \right]^{4-} > \text{PO}_{4}^{3-} > \text{SO}_{4}^{2-} > \text{Cl}^{-}$$

Ans. D

- **Sol.** flocculating power α charge on the ion
- 45. Gold sol is not a:
 - (A) Macromolecular colloid
 - (B) Lyophobic colloid
 - (C) Multimolecular colloid
 - (D) Negatively charged colloid

Ans. A

- Sol. Conceptual
- 46. The *incorrect* statement about Hall -Heroult process is :
 - (A) Carbon anode is oxidised to CO and CO₂
 - (B) Na₃AlF₆ helps to decrease the melting point of the electrolyte
 - (C) CaF₂ helps to increase the conductivity of the electrolyte
 - (D) Oxidation state of oxygen changes in the overall cell reaction

Ans. D

Sol. Conceptual

- 47. Select the correct statement:
 - (A) Roasting involves heating the ore in the absence of air
 - (B) Calcination involves heating the ore above its melting point
 - (C) Smelting involves heating the ore with suitable reducing agent and flux below it melting point
 - (D) Calcination of calcium carbonate is endothermic

Ans. D

- Sol. Conceptual
- 48. NO_2 gas is:
 - (A) Colourless, neutral
 - (B) Colourless, acidic
 - (C) Brown, acidic
 - (D) Brown neutral

Ans. C

- Sol. Conceptual
- 49. Identify the incorrect statement from the following:
 - (A) Oxides of nitrogen in the atmosphere can cause depletion of the ozone layer
 - (B) Ozone absorbs the intense ultraviolet radiation of Sun
 - (C) Depletion of ozone layer is because of its chemical reactions with chlorofluoro alkanes
 - (D) Ozone absorbs infrared radiation

Ans. I

- Sol. Ozone absorbs U.V radiation
- 50. The correct decreasing order of boiling point of hydrogen halides is:
 - (A) HF > HC1 > HBr > HI
 - (B) HI > HBr > HCl > HF
 - (C) HF > HI > HBr > HCl
 - (D) HI > HF > HBr > HC1

Ans. C

- Sol. Conceptual
- 51. The synthetically produced radioactive noble gas by the collision of $^{249}_{98}$ Cf with $^{48}_{20}$ Ca is :
 - (A) Radon
 - (B) Radium
 - (C) Oganesson
 - (D) Xenon

Ans. C

Sol. Oganesson Z=118 $^{249}_{98}$ Cf + $^{48}_{20}$ Ca $\rightarrow ^{294}_{118}$ Og + $^{1}_{0}$ n



- 52. The transition element $(\approx 5\%)$ present with lanthanoid metal in Misch metal is :
 - (A) Mg
- (B) Fe
- (C) Zn
- (D) Co

Ans. B

- **Sol.** 95% lanthanoid metal and $\approx 5\%$ iron
- 53. Match the following:
 - I. Zn²⁺
- i. d⁸ configuration
- II. Cu²⁺
- ii. Colourless
- III. Ni²⁺
- iii. $\mu = 1.73 \text{ BM}$
- Codes:
 - I
- (A) i
- II III ii iii
- (B) i
-
- (B) ii
- iii i
 i iii
- (C) ii (D) i
- iii
- Ans. B
- **Sol.** Zn⁺² no unpaired electrons

$$Cu^{+2} - \mu = 1.73 \text{ BM}$$

Ni⁺² - d⁸ configuration

- 54. Which of the following statements related to lanthanoids is *incorrect*?
 - (A) Lanthanoids are silvery white soft metals.
 - (B) Samarium shows +2 oxidation state.
 - (C) Ce⁺⁴ solutions are widely used as oxidising agents in titrimetric analysis.
 - (D) Colour of Lanthanoid ion in solution is due to d-d transition.

Ans. D

- **Sol.** Colour of Lanthanoid ion in solution is due to f-f transition
- 55. A metalloid is:
 - (A) Bi
 - (B) Sb
 - (C) P
 - (D) Se

Ans. B and D

Sol. Sb and Se are metalloids.

(According to NCERT of 15th and 16th group)

- 56. A pair of isoelectronic species having bond order of one is :
 - (A) N₂, CO
 - (B) N₂, NO⁺
 - (C) O_2^{2-} , F_2
 - (D) CO, NO+


Ans. C

Sol. O_2^{2-} , $F_2 = 18$ electrons each

- 57. Identify the **wrong** relation for real gases:
 - (A) $Z = \frac{V_{ideal}}{V_{real}}$
 - (B) $p_{ideal} = p_{real} + \frac{an^2}{V^2}$
 - (C) $V_{real} = V_{ideal} nb$
 - (D) $\left(p + \frac{a}{V^2}\right)(V b) = RT$

Ans. A

- **Sol.** Compressibility factor $(Z) = \frac{V_{real}}{V_{ideal}}$
- 58. From the diagram



 Δ_r H for the reaction C \rightarrow A is:

- (A) + 35 J
- (B) -15 J
- (C) -35 J
- (D) + 15 J

Ans. C

Sol. A $\xrightarrow{\Delta H_1}$ $2B \xrightarrow{\Delta H_2}$ C

$$\Rightarrow A \rightarrow C$$

$$\Delta H = \Delta H_1 + \Delta H_2 = 35J$$

Then
$$C \rightarrow A$$

$$\Delta H = -35J$$

- 59. For which one of the following mixtures is composition uniform throughout?
 - (A) Sand and water
 - (B) Grains and pulses with stone
 - (C) Mixture of oil and water
 - (D) Dilute aqueous solution of sugar

Ans. D

- **Sol.** Dilute aqueous solution of sugar is a homogeneous solution.
- 60. The energy associated with first orbit of He⁺ is:
 - (A) 0 J
 - (B) -8.72 x 10⁻¹⁸ J
 - (C) -4.58 x 10⁻¹⁸ J
 - (D) -0.545 x 10⁻¹⁸ J

Ans. B

Sol.
$$E_n = -2.18 \times 10^{-18} \times \frac{Z^2}{n^2}$$

$$Z = 2, n = 1$$

$$E = -2.18 \times 10^{-18} \times \frac{4}{1} = -8.72 \times 10^{-18} J$$



