

13. The interval in which the function $f(x) = x^3 - 6x^2 + 9x + 10$ is increasing in
- 1) $(-\infty, 1) \cup (3, \infty)$ 2) $(-\infty, 1] \cup [3, \infty)$
3) $[1, 3]$ 4) $(-\infty, -1] \cup [3, \infty)$

ANS : 2

14. The sides of an equilateral triangle are increasing at the rate of 4 cm/sec. The rate at which its area is increasing. When the side is 14 cm
- 1) $10\sqrt{3} \text{ cm}^2 / \text{sec}$ 2) $14\sqrt{3} \text{ cm}^2 / \text{sec}$
3) $42 \text{ cm}^2 / \text{sec}$ 4) $14 \text{ cm}^2 / \text{sec}$

ANS : GRACE ($28\sqrt{3} \text{ cm}^2 / \text{sec}$)

15. The value of $\sqrt{24.99}$ is
- 1) 4.999 2) 4.899 3) 5.0014) 4.897

ANS : 1

16. If $|3x-5| \leq 2$ then
- 1) $-1 \leq x \leq \frac{7}{3}$ 2) $1 \leq x \leq \frac{7}{3}$
3) $1 \leq x \leq \frac{9}{3}$ 4) $-1 \leq x \leq \frac{9}{3}$

ANS : 2

17. A random variable 'X' has the following probability distribution

X	1	2	3	4	5	6	7
P(X)	k-1	3k	K	3k	3k ²	k ²	k ² +k

Then the value of k is

- 1) $\frac{1}{5}$ 2) -2 3) $\frac{2}{7}$ 4) $\frac{1}{10}$

ANS : 1

18. If A and B are two events of a sample space S such that $P(A)=0.2$, $P(B)=0.6$ and $P(A|B)=0.5$ then $P(A'|B)$
- 1) $\frac{3}{10}$ 2) $\frac{2}{3}$ 3) $\frac{1}{2}$ 4) $\frac{1}{3}$

ANS : 3

19. If 'X' has a binomial distribution with parameter $n=6$, p and $P(X=2)=12$, $P(X=3)=5$ then $P=$
- 1) $\frac{5}{12}$ 2) $\frac{16}{21}$ 3) $\frac{1}{2}$ 4) $\frac{5}{6}$

ANS : GRACE ($P = \frac{5}{21}$)

20. A man speaks truth 2 out of 3 times. He picks one of the natural number in the set $S = \{1, 2, 3, 4, 5, 6, 7\}$ and reports that it is even. The probability that it is actually even is
- 1) $\frac{2}{5}$ 2) $\frac{1}{5}$ 3) $\frac{1}{10}$ 4) $\frac{3}{5}$

ANS : 4

21. $\int_{-3}^3 \cot^{-1} x \, dx =$
- 1) 3π 2) 0 3) 6π 4) 3

ANS : 1

22. $\int \frac{1}{\sqrt{x} + x\sqrt{x}} \, dx =$
- 1) $2 \log(\sqrt{x} + 1) + C$ 2) $\frac{1}{2} \tan^{-1} \sqrt{x} + C$
3) $\tan^{-1} \sqrt{x} + C$ 4) $2 \tan^{-1} \sqrt{x} + C$

ANS : 4

23. $\int \frac{2x-1}{(x-1)(x+2)(x-3)} \, dx = A \log|x-1| + B \log|x+2| + C \log|x-3| + K$, then A, B, C are respectively
- 1) $\frac{-1}{6}, \frac{1}{3}, \frac{-1}{2}$ 2) $\frac{1}{6}, \frac{1}{3}, \frac{1}{5}$
3) $\frac{1}{6}, \frac{-1}{3}, \frac{1}{3}$ 4) $\frac{-1}{6}, \frac{-1}{3}, \frac{1}{2}$

ANS : 4

24. $\int_0^2 [x^2] \, dx =$
- 1) $5 - \sqrt{2} - \sqrt{3}$ 2) $5 + \sqrt{2} - \sqrt{3}$
3) $5 - \sqrt{2} + \sqrt{3}$ 4) $-5 - \sqrt{2} - \sqrt{3}$

ANS : 1

25. $\int_0^1 \sqrt{\frac{1+x}{1-x}} dx =$

- 1) $\frac{\pi}{2} - 1$ 2) $\frac{\pi}{2} + 1$ 3) $\frac{\pi}{2}$ 4) $\frac{1}{2}$

ANS : 2

26. If $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: [0, \infty) \rightarrow \mathbb{R}$ is defined by $f(x) = x^2$ and $g(x) = \sqrt{x}$. Which one of the following is not true?

- 1) $\text{gof}(4) = 4$ 2) $\text{fog}(-4) = 4$
3) $\text{fog}(2) = 2$ 4) $\text{gof}(-2) = 2$

ANS : 2

27. If $A = \{x | x \in \mathbb{N}, x \leq 5\}$, $B = \{x | x \in \mathbb{Z}, x^2 - 5x + 6 = 0\}$, then the number of onto functions from A to B is

- 1) 2 2) 23 3) 30 4) 32

ANS : 3

28. On the set of positive rationals, a binary operation $*$ is defined by $a * b = \frac{2ab}{5}$. If

$2 * x = 3^{-1}$ then $x =$

- 1) $\frac{1}{6}$ 2) $\frac{5}{12}$ 3) $\frac{2}{5}$ 4) $\frac{125}{48}$

ANS : 4

29. $\cos \cos \left[2 \sin^{-1} \frac{3}{4} + \cos^{-1} \frac{3}{4} \right] =$

- 1) $\frac{-3}{4}$ 2) $\frac{3}{4}$
3) $\frac{3}{5}$ 4) does not exist

ANS : 1

30. If $a + \frac{\pi}{2} < 2 \tan^{-1} x + 3 \cot^{-1} x < b$ then 'a' and 'b' are respectively.

- 1) 0 and π 2) $\frac{\pi}{2}$ and 2π
3) $\frac{\pi}{2}$ and 2π 4) $\frac{-\pi}{2}$ and $\frac{\pi}{2}$

ANS : 2

31. If U is the universal set with 10 elements; A and B are two sets such that $n(A) = 50$, $n(B) = 60$, $n(A \cap B) = 20$ then $n(A' \cap B') =$

- 1) 40 2) 20 3) 90 4) 10

ANS : 4

32. The domain of the function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \sqrt{x^2 - 7x + 12}$ is

- 1) $(-\infty, 3] \cup [4, \infty)$ 2) $(-\infty, 3] \cup (4, \infty)$
3) $(-\infty, 3] \cap [4, \infty)$ 4) (3, 4)

ANS : 1

33. If $\cos x = |\sin x|$ then, the general solution is

- 1) $x = n\pi \pm \frac{\pi}{4}$, $n \in \mathbb{Z}$
2) $x = 2n\pi \pm \frac{\pi}{4}$, $n \in \mathbb{Z}$
3) $x = n\pi \pm (-1)^n \frac{\pi}{4}$, $n \in \mathbb{Z}$
4) $x = (2n+1)\pi \pm \frac{\pi}{4}$, $n \in \mathbb{Z}$

ANS : 1

34. $\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ =$

- 1) 2 2) 3 3) 4 4) 1

ANS : 3

35. If $P(n): 2^n < n!$ then the smallest positive integer for which P(n) is true, is

- 1) 2 2) 3 3) 4 4) 5

ANS : 3

36. Foot of the perpendicular drawn from the point (1, 3, 4) to the plane $2x - y + z + 3 = 0$ is

- 1) (-1, 4, 3) 2) (0, -4, -7)
3) (1, 2, -3) 4) (-3, 5, 2)

ANS : 1

37. Acute angle between the line $\frac{x-5}{2} = \frac{y+1}{-1} = \frac{z+4}{1}$ and the plane $3x - 4y - z + 5 = 0$ is

- 1) $\cos^{-1} \left(\frac{9}{\sqrt{364}} \right)$ 2) $\sin^{-1} \left(\frac{9}{\sqrt{364}} \right)$
3) $\cos^{-1} \left(\frac{5}{2\sqrt{13}} \right)$ 4) $\sin^{-1} \left(\frac{5}{2\sqrt{13}} \right)$

ANS : 3

38. The distance of the point (1, 1, 1) from the line $\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-3}{2}$ is


- 1) $\frac{2\sqrt{3}}{5}$ 2) $\frac{2\sqrt{5}}{3}$ 3) $\frac{\sqrt{5}}{3}$ 4) $\frac{20}{3}$

ANS : 2

39. XY-plane divides the line joining the points A(2, 3, -5) and B(-1, -2, -3) in the ratio
 1) 2:1 internally 2) 3:2 externally
 3) 5:3 internally 4) 5:3 externally

ANS : 4

40. The shaded region in the figure is the solution set of the inequations



- 1) $4x+5y \geq 20, 3x+10y \leq 30, x \leq 6, x, y \geq 0$
 2) $4x+5y \geq 20, 3x+10y \leq 30, x \geq 6, x, y \geq 0$
 3) $4x+5y \leq 20, 3x+10y \leq 30, x \leq 6, x, y \geq 0$
 4) $4x+5y \leq 20, 3x+10y \leq 30, x \geq 6, x, y \geq 0$

ANS : 1

41. The inverse of the matrix $\begin{bmatrix} 2 & 5 & 0 \\ 0 & 1 & 1 \\ -1 & 0 & 3 \end{bmatrix}$ is

- 1) $\begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$ 2) $\begin{bmatrix} 3 & -5 & 5 \\ -1 & -6 & -2 \\ 1 & -5 & 2 \end{bmatrix}$
 3) $\begin{bmatrix} 3 & -15 & 5 \\ -1 & 6 & -2 \\ 1 & -5 & 2 \end{bmatrix}$ 4) $\begin{bmatrix} 3 & -15 & 5 \\ -1 & 6 & -2 \\ 1 & -5 & -2 \end{bmatrix}$

ANS : 3

42. If P and Q are symmetric matrices of the same order then PQ-QP is
 1) identify matrix
 2) symmetric matrix
 3) zero matrix
 4) skew symmetric matrix

ANS : 4

43. If $3A + 4B' = \begin{bmatrix} 7 & -10 & 17 \\ 0 & 6 & 31 \end{bmatrix}$ and
 $2B - 3A' = \begin{bmatrix} -1 & 18 \\ 4 & 0 \\ -5 & -7 \end{bmatrix}$ then B =

- 1) $\begin{bmatrix} 1 & 3 \\ -1 & 1 \\ 2 & 4 \end{bmatrix}$ 2) $\begin{bmatrix} 1 & -3 \\ -1 & 1 \\ 2 & 4 \end{bmatrix}$
 3) $\begin{bmatrix} -1 & -18 \\ 4 & -16 \\ -5 & -7 \end{bmatrix}$ 4) $\begin{bmatrix} 1 & 3 \\ -1 & 1 \\ 2 & -4 \end{bmatrix}$

ANS : 1

44. If $A = \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix}, B = \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}$, then $|ABB'| =$
 1) 50 2) -250 3) 100 4) 250

ANS : 2

45. If the value of a third order determinant is 16, then the value of the determinant formed by replacing each of its elements by its cofactor is
 1) 96 2) 48 3) 256 4) 16

ANS : 3

46. The eccentricity of the ellipse $9x^2 + 25y^2 = 225$ is
 1) $\frac{4}{5}$ 2) $\frac{3}{5}$ 3) $\frac{3}{4}$ 4) $\frac{9}{16}$

ANS : 1

47. $\sum_{r=1}^n (2r-1) = x$ then

$$\lim_{n \rightarrow \infty} \left[\frac{1^3}{x^2} + \frac{2^3}{x^2} + \frac{3^3}{x^2} + \dots + \frac{n^3}{x^2} \right] =$$

1) $\frac{1}{2}$ 2) $\frac{1}{4}$ 3) 1 4) 4

ANS : 2

48. The negation of the statement "All continuous functions are differentiable."
 1) All continuous functions are not differentiable
 2) Some continuous functions are differentiable
 3) Some continuous functions are not differentiable
 4) All differentiable functions are continuous

ANS : 3

49. Mean and standard deviation of 100 items are 50 and 4 respectively. The sum of all squares of the items is
1) 251600 2) 256100 3) 266000 4) 261600

ANS : 1

50. Two letters are chosen from the letters of the word 'EQUATIONS'. The probability that one is vowel and the other is consonant is

- 1) $\frac{8}{9}$ 2) $\frac{4}{9}$ 3) $\frac{3}{9}$ 4) $\frac{5}{9}$

ANS : 4

51. $\int x^3 \sin 3x \, dx =$

- 1) $-\frac{x^3 \cos 3x}{3} - \frac{x^2 \sin 3x}{3} + \frac{2x \cos 3x}{9} - \frac{2 \sin 3x}{27} + C$
2) $\frac{x^3 \cos 3x}{3} + \frac{x^2 \sin 3x}{3} - \frac{2x \cos 3x}{9} - \frac{2 \sin 3x}{27} + C$
3) $-\frac{x^3 \cos 3x}{3} + \frac{x^2 \sin 3x}{3} + \frac{2x \cos 3x}{9} - \frac{2 \sin 3x}{27} + C$
4) $-\frac{x^3 \cos 3x}{3} + \frac{x^2 \sin 3x}{3} - \frac{2x \cos 3x}{9} - \frac{2 \sin 3x}{27} + C$

ANS : 4

52. The area of the region above X-axis included between the parabola $y^2=x$ and the circle $x^2+y^2=2x$ in square units is

- 1) $\frac{\pi}{4} - \frac{3}{2}$ 2) $\frac{3}{2} - \frac{\pi}{4}$ 3) $\frac{2}{3} - \frac{\pi}{4}$ 4) $\frac{\pi}{4} - \frac{2}{3}$

ANS : 4

53. The area of the region bounded by Y-axis, $y=\cos x$ and $y=\sin x$; $0 \leq x \leq \frac{\pi}{2}$ is

- 1) $\sqrt{2} - 1$ Sq. units 2) $\sqrt{2}$ Sq. units
3) $\sqrt{2} + 1$ Sq. units 4) $2 - \sqrt{2}$ Sq. units

ANS : 1

54. The integrating factor of the differential equation $(2x+3y^2) dy=y dx(y>0)$ is

- 1) $e^{\frac{1}{y}}$ 2) $-\frac{1}{y^2}$ 3) $\frac{1}{x}$ 4) $\frac{1}{y^2}$

ANS : 4

55. The equation of the curve passing through the point (1, 1) such that the slope of the tangent at any point (x, y) is equal to the product of its co-ordinates is

- 1) $2 \log x = y^2 - 1$ 2) $2 \log y = x^2 + 1$
3) $2 \log y = x^2 - 1$ 4) $2 \log x = y^2 + 1$

ANS : 1

56. The constant term in the expansion of

$$\left| \begin{matrix} 3x+1 & 2x-1 & x+2 \\ 5x-1 & 3x+2 & x+1 \\ 7x-2 & 3x+1 & 4x-1 \end{matrix} \right| \text{ is}$$

- 1) 0 2) 2 3) -10 4) 6

ANS : 4

57. If $[x]$ represents the greatest integer function

and $f(x) = x - [x] - \cos x$ then $f'\left(\frac{\pi}{2}\right) =$

- 1) 0 2) 1 3) 2
4) does not exist

ANS : 3

58. If $f(x) = \begin{cases} \frac{\sin 3x}{e^{2x}-1} & ; x \neq 0 \\ k-2 & ; x = 0 \end{cases}$ is

continuous at $x=0$, then $k=$

- 1) $\frac{3}{2}$ 2) $\frac{9}{5}$ 3) $\frac{1}{2}$ 4) $\frac{2}{3}$

ANS : GRACE $\left(\frac{7}{2}\right)$

59. if $f(x) = \sin^{-1} \left[\frac{2^{x+1}}{1+4^x} \right]$, then $f'(0) =$

- 1) $2 \log 2$ 2) $\log 2$ 3) $\frac{2 \log 2}{5}$ 4) $\frac{4 \log 2}{5}$

ANS : 2

60. If $x = a \sec^2 \theta$, $y = a \tan^2 \theta$ then $\frac{d^2 y}{dx^2} =$

- 1) $2a$ 2) 1 3) 0 4) 4

ANS : 3